VOL. 631 NOS. 1+2 JOURNAL OF CHROMATOGRAPHY FEBRUARY 12, 1993

CONTENTS

16TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, BALTIMORE, MD, JUNE 14–19, 1992, PART I

COLUMN LIQUID CHROMATOGRAPHY

Theory and general

Chromatographic method verification by means of three-dimensional, multivariate visualization
by F. O. Geiser (Glen Mills, PA, USA), C. Golt and L. Kung, Jr. (Newark, DE, USA), J. D. Justice (Palo Alto, CA, USA) and B. L. Brown (Orem, UT, USA) 1

Faster quantitative evaluation of high-performance liquid chromatography–ultraviolet diode-array data by multicomponent analysis
by J.-L. Excoffier, M. Joseph, J.J. Robinson and T. L. Sheehan (Walnut Creek, CA, USA) 15

Deuterium nuclear magnetic resonance spectroscopy as a probe for reversed-phase liquid chromatographic bonded phase solvation: methanol and acetonitrile mobile phase components
by D. M. Bliesner and K. B. Sentell (Burlington, VT, USA) 23

Thermodynamics of the adsorption of Tröger's base enantiomers from ethanol on cellulose triacetate
by A. Seidel-Morgenstern and G. Guiochon (Knoxville and Oak Ridge, TN, USA) 37

New chromatographic hydrophobicity index (φ_0) based on the slope and the intercept of the log k' *versus* organic phase concentration plot
by K. Valkó and P. Slégel (London, UK) 49

Accurate determination of log k'_w in reversed-phase liquid chromatography. Implications for quantitative structure–retention relationships
by M.-M. Hsieh and J. G. Dorsey (Cincinnati, OH, USA) 63

Materials and techniques

Characterization of a Chiral-AGP capillary column coupled to a micro sample-enrichment system with UV and electrospray mass spectrometric detection
by J. Hermansson and I. Hermansson (Norsborg, Sweden) and J. Nordin (Huddinge, Sweden) 79

Reversed-phase liquid chromatography with microspherical octadecyl-zirconia bonded stationary phases
by J. Yu and Z. El Rassi (Stillwater, OK, USA) 91

Fimbriated stationary phases for proteins
by L. Varady (Cambridge, MA, USA), N. Mu (West Lafayette, IN, USA), Y.-B. Yang (Hesperia, CA, USA), S. E. Cook (Wilmington, DE, USA), N. Afeyan (Cambridge, MA, USA) and F. E. Regnier (West Lafayette, IN, USA) 107

High-performance liquid chromatography of amino acids, peptides and proteins. CXXIV. Physical characterisation of fluidized-bed behaviour of chromatographic packing materials
by G. Dasari, I. Prince and M. T. W. Hearn (Clayton, Australia) 115

Selectivity of organic solvents in micellar liquid chromatography of amino acids and peptides
by A. S. Kord and M. G. Khaledi (Raleigh, NC, USA) 125

Axial illumination of fused-silica capillaries. Investigation of fluorescence and refractive index detection
by A. A. Abbas and D. C. Shelly (Lubbock, TX, USA) 133

Peak tracking with a neural network for spectral recognition
by P. M. J. Coenegracht, H. J. Metting, E. M. van Loo, G. J. Snoeijer and D. A. Doornbos (Groningen, Netherlands) 145

Primary contribution of the injector to carryover of a trace analyte in high-performance liquid chromatography
by M. Saha and R. W. Giese (Boston, MA, USA) 161

Applications: chiral separations

N-Arylcarbamoyl derivatives of amino acids as chiral stationary phases for optical resolution by high-performance liquid chromatography
by M.-H. Yang and J.-Y. Lin (Taipei, Taiwan) 165

Analytical and preparative high-performance liquid chromatographic separation of thienopyran enantiomers
by C. J. Shaw, P. J. Sanfilippo, J. J. McNally, S. A. Park and J. B. Press (Raritan, NJ, USA) 173

Enantiomer separation by high-performance liquid chromatography with copper(II) complexes of Schiff bases as chiral stationary phases
by N. Ôi, H. Kitahara and F. Aoki (Osaka, Japan) 177

Retention and enantioselective properties of racemic compounds on modified ovomucoid columns. II. Reaction with glyceraldehyde, formaldehyde and glutaric anhydride
by J. Haginaka, T. Murashima and C. Seyama (Hyogo, Japan) and H. Fujima and H. Wada (Kyoto, Japan) 183

Direct chiral separation of almokalant on Chiralcel OD and Chiralpak AD for liquid chromatographic assay of biological samples
by K. Balmér, P.-O. Lagerström, S. Larsson and B.-A. Persson (Mölndal, Sweden) 191

Rapid direct resolution of the stereoisomers of all-*trans* astaxanthin on a Pirkle covalent L-leucine column
by S. A. Turujman (Washington, DC, USA) 197

Applications: pharmaceuticals

Determination of ephedrines in urine by high-performance liquid chromatography
by C. Imaz, D. Carreras, R. Navajas, C. Rodriguez, A. F. Rodriguez, J. Maynar and R. Cortes (Madrid, Spain) . . . 201

Porous graphitized carbon and octadecylsilica columns in the separation of some monoamine oxidase inhibitory drugs
by E. Forgács, K. Valkó, T. Cserháti and K. Magyar (Budapest, Hungary) 207

Liquid chromatography–mass spectrometry for the determination of medetomidine and other anaesthetics in plasma
by H. Kanazawa, Y. Nagata, Y. Matsushima, N. Takai, H. Uchiyama, R. Nishimura and A. Takeuchi (Tokyo, Japan) 215

Reversed-phase high-performance liquid chromatography of the cardiac glycoside LNF-209 with refractive index detection
by P. H. Zoutendam, D. L. Berry and D. W. Carkuff (Norwich, NY, USA) 221

Determination of oxiracetam in human plasma by reversed-phase high-performance liquid chromatography with fluorimetric detection
by R. C. Simpson, V. K. Boppana, B. Y.-H. Hwang and G. R. Rhodes (King of Prussia, PA, USA) 227

Column-switching high-performance liquid chromatographic method for the determination of SK&F 106203 in human plasma after fluorescence derivatization with 9-anthryldiazomethane
by C. Miller-Stein, B. Y.-H. Kwang, G. R. Rhodes and V. K. Boppana (King of Prussia, PA, USA) 233

Determination of baicalin and puerarin in traditional Chinese medicinal preparations by high-performance liquid chromatography
by K.-C. Wen, C.-Y. Huang and F.-L. Lu (Taipei, Taiwan) 241

Normal-phase high-performance liquid chromatographic determination of epristeride, a prostatic steroid 5α-reductase enzyme inhibitor, in human plasma
by V. K. Boppana, C. Miller-Stein and G. R. Rhodes (King of Prussia, PA, USA) 251

Rapid, sensitive high-performance liquid chromatographic method for the quantification of promethazine in human serum with electrochemical detection
by A. R. Fox and D. A. McLoughlin (Baltimore, MD, USA) 255

Applications: proteins and their constituents

High-performance liquid chromatography of human glycoprotein hormones
by M. A. Chlenov, E. I. Kandyba, L. V. Nagornaya, I. L. Orlova and Y. V. Volgin (Moscow, Russian Federation) . 261

Immunoaffinity chromatography of recombinant *Amb a* I in the presence of a denaturing agent
by K. M. Keating, B. L. Rogers, L. Weber, J. P. Morgenstern, D. G. Klapper and M. Kuo (Cambridge, MA and Chapel Hill, NC, USA) 269

Separation of interleukins by a preparative chromatofocusing-like method
by S. P. Monkarsh, E. A. Russoman and S. K. Roy (Nutley, NJ, USA) 277

Kinetic analysis of biotinylation of specific residues of peptides by high-performance liquid chromatography
by A. Korosky, B. T. Miller and S. L. Knock (Galveston, TX, USA) 281